

Escherichia coli DH5 α

Multiporator[®] / Electroporator 2510

Transformation Protocol

Protocol No. 4308 915.512 – 04/2002

Microorganism	<i>Escherichia coli</i> DH5 α
Cell type	Bacteria, gram negative
Molecules injected	Plasmid DNA (pUC 19)
Growth medium	LB medium
Washing solution	Sterile, ice-cold water; (10% glycerol)
Electroporation solution	Sterile, ice-cold water; (10% glycerol)
Outgrowth medium	SOC medium (without antibiotics)
Cuvette	1 mm gap width
Reference	Eppendorf AG • Application Hotline • D-22331 Hamburg Phone +49 180 3 666 789 • Fax +49 40 53990 125 • e-mail: application-hotline@eppendorf.de Adapted from: High Efficiency Transformation by Electroporation • Short Protocols in Molecular Biology Second Edition • Green Publishing Association and John Wiley & Sons, New York • 1-22 – 1-23

Making electrocompetent cells:

1. Inoculate 500 ml LB medium with 2.5 ml of a fresh overnight culture of *E. coli* DH5 α . Grow at 37 °C with shaking to an O.D.₆₀₀ of 0.5 to 0.6.
2. Chill cells on ice for 15 minutes and transfer to a prechilled centrifuge bottle. Harvest by centrifugation (20 minutes, 5,000 x g, 2-4 °C). Resuspend pellet in 5 ml ice-cold water. Keep the cells cold during the entire procedure.
3. Wash twice with the original culture volume of ice-cold water. Centrifuge as above. Resuspend pellet by swirling in remaining liquid.
- 4a If using the cells immediately, place suspension in a prechilled tube and centrifuge (10 minutes, 5,000 x g, 2-4 °C). Resuspend the cells in ice-cold water to a final concentration of approximately 2×10^{11} cells/ml. Aliquote 40-300 μ l cells into prechilled centrifuge tubes.
- 4b If freezing the cells for later use, add 40 ml of ice-cold 10% glycerol, mix and centrifuge for 10 minutes, at 5,000 x g and 2-4 °C. Resuspend the cells in ice-cold 10% glycerol to a final concentration of approximately, 2×10^{11} cells/ml. Aliquote 40-300 μ l cells into prechilled centrifuge tubes. quick freeze on dry ice and store at -80 °C.

Electroporation of cells:

1. Add 1 μ l DNA (10 pg in water) to tubes containing 40 μ l electrocompetent cells. Homogenize by gently mixing with pipette several times.
2. Transfer mixture to a prechilled cuvette. Wipe moisture from the cuvette and insert the cuvette into the device.
3. Electroporation:

Mode	Prokaryotes "O"
Voltage (V)	1,700 V
Time constant (τ)	5 ms

4. Immediately add 1 ml SOC medium and transfer to a sterile culture tube with a pasteur pipette. Incubate 30-60 minutes with moderate shaking at 37 °C.
5. Plate on LB plates containing the appropriate selection chemical.

Expected results:

Transformation efficiency up to 3×10^9 transformants/ μ g of DNA.

Brinkmann Instruments

Support and Services Directory

Contact Information

United States

Canada



Business Hours:

8:30 a.m. to 6:00 p.m. EST

8:30 a.m. to 6:00 p.m. EST



Phone:

800-645-3050
516-334-7500

800-263-8715
905-826-5525

Fax:

516-334-7506

905-826-5424



Address:

One Cantiague Road
P.O. Box 1019
Westbury, NY 11590-0207

6670 Campobello Road
Mississauga, ONT L5N 2L8



Website:
E-mail:

www.brinkmann.com
info@brinkmann.com

www.brinkmann.com
canada@brinkmann.com



Customer Support:

800-645-3050, menu option 2
custserv@brinkmann.com

800-263-8715, ext. 237
custserv@brinkmann.com



Repair:

800-645-3050, ext. 2404
techserv@brinkmann.com

800-263-8715, ext. 232
techserv@brinkmann.com



BioSystems Lab:

800-645-3050, ext. 2258
bioapps@brinkmann.com

516-334-7500, ext. 2258 (U.S.)
bioapps@brinkmann.com

Analytical Systems Lab:

800-645-3050, ext. 2421
apps@brinkmann.com

516-334-7500, ext. 2421 (U.S.)
apps@brinkmann.com

brinkmann
— an eppendorf company —